

# ELECTROWEAK PHASE TRANSITION AND BARYOGENESIS IN THE MINIMAL SUPERSYMMETRIC STANDARD MODEL

J.M. CLINE

*McGill University, Department of Physics, Montréal, Québec H3A 2T8, Canada*

I describe work done in collaboration with M. Joyce and K. Kainulainen on (1) the strength of the electroweak phase transition in the MSSM and (2) the mechanism for producing the baryon asymmetry during the phase transition. In the former we compare the effective potential and dimensional reduction methods for describing the phase transition and search the parameter space of the MSSM for those values where it is strong enough. In the latter we give a systematic computation of the baryon asymmetry due the CP-violating force acting on charginos in the vicinity of the bubble wall. We find that a light right-handed stop, a light Higgs boson, and a large phase in the  $\mu$  parameter, are the main necessary ingredients for producing the baryon asymmetry.

## 1 Experimentally testable baryogenesis?

A major triumph of astroparticle physics is our understanding of the light element abundances through big bang nucleosynthesis. Equally exciting is the prospect of putting the measured baryon asymmetry of the universe,  $\eta = n_B/n_\gamma \cong 3 \times 10^{-10}$ , on a similar footing. Unfortunately, there is much less agreement about the mechanism of baryogenesis than there is for nucleosynthesis. A search of the SPIRES database shows about 400 papers on the subject so far, 350 of which (my rough estimate) describe different ways of generating the baryon asymmetry. Which one of them, if any, is correct? Of course it is possible that the asymmetry was generated at such high temperatures that direct confirmation of the new physics needed will never be achieved in the laboratory. In this case baryogenesis should forever be relegated to the realm of untestable speculation, which is not very interesting from the point of view of phenomenology.

More interesting are the scenarios which can be falsified experimentally before we retire. Electroweak baryogenesis is arguably the most attractive such possibility at present. It needs new physics at the 100 GeV energy scale, of a kind that is even now being probed by the LEP 2 experiment. It thus holds out the hope that it could be ruled out in the near future, reducing the number of candidate baryogenesis mechanisms from 350 to 349: this is progress! On the other hand, if it is not ruled out we may be on the way to a truly quantitative theory of the origin of baryonic matter, with consequences more profound than nucleosynthesis for our understanding of particle physics and the early universe.




Figure 1: Expanding bubble during the electroweak phase transition. Left-handed fermions reflect into right-handed ones and vice versa, but with different probabilities.

The basic picture of how electroweak baryogenesis could work was established by Cohen, Kaplan and Nelson<sup>1</sup> in 1992. The electroweak phase transition is supposed to be first order, meaning that at some critical temperature  $T \sim 100$  GeV, the bubbles of the broken phase (where the Higgs field VEV is nonzero) begin to nucleate in the high temperature symmetric phase (where  $\langle H \rangle = 0$ ). The bubbles expand, collide and eventually fill up the universe with the broken phase. Before doing so however, particles in the high- $T$  plasma will be partially reflected when encountering the bubble walls, as can be understood from the fact that they gain mass from the nonzero Higgs VEV as they enter the bubbles. These interactions with the bubble walls can be CP violating. For example, left-handed top quarks may reflect from the bubble walls more strongly than right-handed top quarks, leading to a buildup of chirality in front of the expanding bubble wall.

Enter the sphaleron, the electroweak field configurations which violate baryon number in the Standard Model due to the fact that the baryon current is anomalous in the presence of background  $W$  boson fields. Each sphaleron interaction creates or destroys 9 left-handed quarks and 3 left-handed leptons in such a way as to violate  $B$  (baryon number) and  $L$  (lepton number) by 3 units, while preserving  $B - L$ . Sphalerons, “seeing” the excess of chirality in front of the wall, will tend to drive it to zero by converting excess left-handed quarks into their antiparticles. However this in turn creates a baryon asymmetry. Eventually the excess baryons in front of the wall are overtaken by the wall and fall inside the bubbles, where the sphalerons are much heavier than they are in the symmetric phase. If the sphalerons are sufficiently Boltzmann-suppressed inside the bubbles, the baryon excess is safe from erasure by sphalerons, which would otherwise, given enough time, tend to cause it to relax to zero. (The competing tendency to erase chirality is no longer




Figure 2: Schematic picture of the effective potential of the Higgs field near the critical temperature.

present inside the bubbles because the particles have masses there, and so chirality is not a good quantum number.)

New physics is needed by electroweak baryogenesis for the following reasons. (1) The phase transition should be strongly enough first order to prevent the sphalerons from erasing the baryon asymmetry inside the bubbles, and (2) the CP violation on the wall must be strong enough to generate enough baryons. A natural candidate for the origin of such new physics is the minimal supersymmetric standard model (MSSM), since it is able to fulfill both of these criteria.<sup>2</sup>

## 2 Strength of the phase transition: when is $v_c/T_c > 1$ ?

To avoid the washout of baryons inside the bubbles once they have been created, the sphaleron configuration must be heavy compared to the temperature. In this case the rate of sphaleron interactions inside the bubbles goes like  $\Gamma_{\text{sph}} \sim e^{-\text{const.}(v_c/T_c)}$  where  $v_c$  is the Higgs field VEV (normalized so that  $v = 246$  GeV at  $T = 0$ ) in the bubbles at the critical temperature  $T_c$ , when nucleation takes place (figure 2). This follows from the fact that the sphaleron energy is proportional to  $v_c$ . One can show that baryon preservation requires that  $v_c/T_c$  be greater than 1.0–1.1, the exact number depending on how the potential evolves with temperature. In the standard model it is impossible to satisfy the  $v_c/T_c > 1$  constraint, but in the MSSM it has been shown that the right-handed top squarks, if sufficiently light, can enhance the strength of the phase transition to the required level.

I will review two methods that have been used to compute the critical ratio  $v_c/T_c$ . Traditionally one used the finite-temperature effective potential (EP) at one or two loops.<sup>3</sup> However at high  $T$  the perturbative expansion is no longer controlled by the usual dimensionless couplings, because it is effectively a 3D theory with potential infrared divergences from the light (transverse  $W$  boson)




Figure 3: Distributions of parameters satisfying the baryogenesis phase transition constraint in the MSSM, using dimensional reduction (top line) and the effective potential (bottom line). Masses are in units of 100 GeV.

degrees of freedom whose thermal masses are small. These IR problems can render the predictions of the EP unreliable. Roughly speaking, this happens if the ratio  $\lambda T_c/m \gtrsim 1$ , where  $m$  is the thermal mass of the light particle at  $T_c$  and  $\lambda$  is its quartic coupling to the Higgs boson.

A possible way around this problem is dimensional reduction (DR),<sup>4</sup> where one computes an effective theory by integrating out all *but* these most dangerous infrared contributions, leaving the lattice to take care of them. In DR, an effective three dimensional Lagrangian is obtained which has the form  $\mathcal{L}_{\text{eff}} = |\vec{D}H|^2 + F_{ij}F_{ij}/4g_3^2 - \mu_3^2|H|^2 + \lambda_3|H|^4$ , where  $H$  is the light linear combination of the two Higgs doublets of the MSSM. It has been found<sup>5</sup> that the limit  $v_c/T_c > 1$  corresponds to  $x_c \equiv \lambda_3/g_3^2 < 0.044$ . Thus one need only compute  $x_c$  in terms of the parameters of the MSSM to find which ones are consistent with baryogenesis.

We have searched the MSSM parameter space using both the EP and DR at one loop to compare the two approaches,<sup>6</sup> and found that they are roughly compatible, although DR gives a somewhat bigger range of allowed parameters, as shown in figure 1. The most sensitive parameters along with their predicted values are  $\tan\beta \equiv \langle H_2 \rangle / \langle H_1 \rangle = 2^{+1.5}_{-0.5}$ ,  $m_{h^0} < 85$  GeV, a light stop with  $m_{\tilde{t}_1} = 150 - 200$  GeV, and a correspondingly small right-handed soft-breaking stop mass parameter of  $m_U < 100$  GeV. Negative values of  $m_U^2$  have been advocated for making the transition strong enough, but our results show that although they are helpful, they are not required. Furthermore we find that arbitrarily large values of  $\mu$  are allowed, which is encouraging because  $\mu$  is the major source of CP violation in the model. This is in contrast to previous studies that have emphasized the weakening effect of  $\mu$  on the transition. We will comment upon the phenomenological relevance of these parameter values below.

It should be mentioned that studies of the EP at two loops show a less restrictive range of allowed parameters, more in agreement with DR at one loop.<sup>3</sup> It would be interesting to extend the DR analysis to two loops to see if agreement between the two methods is further improved.

### 3 Origin of the baryon asymmetry—classical force mechanism

Knowing the preferred parameters for safeguarding the baryon asymmetry, we now ask whether it can be generated in sufficient quantity in the first place. At first sight this might appear to be a very daunting problem, because the particles reflecting off the bubble wall are colliding with other particles in the plasma while they are quantum mechanically reflecting from the potential of the bubble wall. This mixture of particle diffusion and quantum reflection might compel us to use the quantum Boltzmann equation, a very unwieldy formalism.

However the bubble wall has a thickness of  $L \sim 15/T$ , while the average particle in the plasma has a DeBroglie wavelength of  $\lambda \sim 0.3/T$ . This means that its interaction with the wall is not really quantum mechanical (such effects are exponentially suppressed when  $\lambda \ll L$ ), but rather can be derived from a classical force  $F$  exerted by the wall on the particles.<sup>7</sup> We are thus justified in using the usual Boltzmann equation,  $\partial f / \partial t - \vec{v} \cdot \vec{\nabla}_x f + \vec{F} \cdot \vec{\nabla}_p f = \mathcal{C}[f]$ . The force  $F$  acts with opposite sign on opposite chiralities, leading to chiral charge separation in front of the wall. This can be derived in a rigorous way, starting from a WKB solution to the particles' equation of motion in the vicinity of the wall.<sup>8</sup>

We find that the most important contribution to the baryon asymmetry is from Winos and Higgsinos reflecting from the wall and subsequently being converted (via top quark Yukawa interactions) into a quark asymmetry that biases the sphalerons. The CP violation appears in the chargino mass matrix through the phase of the  $\mu$  parameter:

$$\begin{pmatrix} \bar{W}_R^+ & \bar{h}_{1,R}^+ \end{pmatrix} \begin{pmatrix} m_2 & gv_2(z)/\sqrt{2} \\ gv_1(z)/\sqrt{2} & |\mu|e^{i\delta} \end{pmatrix} \begin{pmatrix} \tilde{W}_L^+ \\ \tilde{h}_{2,L}^+ \end{pmatrix}.$$

The final answer for the baryon asymmetry depends on  $m_2$  (the Wino mass parameter),  $\mu$ ,  $\tan \beta$ , the Higgs and quark diffusion constants, the wall velocity  $v_w$ , the strong and weak sphaleron rates, the rate of top Yukawa interactions, the wall width, and the critical VEV and temperature. It is also proportional to the underlying CP violation in the  $\mu$  parameter,  $\sin \delta$ . Putting in our best estimates for these quantities (or the ranges allowed by the constraint  $v_c/T_c > 1$ ) we find that typically  $\sin \delta$  must be unity to get a large enough asymmetry, although if  $m_2 \cong \mu$  then it is possible to have  $\sin \delta \sim 0.1$ .




Figure 4: Contours of constant  $\eta_{10} = 10^{10} n_B/n_\gamma$  in the plane of  $m_2$  (the Wino mass) and  $\mu$ , in GeV,  $v_{\text{wall}} = 0.05, 0.1$  and  $0.2$ ,  $\tan \beta = 2$ ,  $\sin \delta = 1$  and wall width  $= 14/T$ .

The dependence of  $\eta_{10} = 10^{10} n_B/n_\gamma$  on  $\mu$  and  $m_2$  is shown for several choices of other parameters in figure 2.

#### 4 Relation to Collider Physics and EDM Searches

We have shown that electroweak baryogenesis in the MSSM works only for rather limited ranges of parameters like  $2.5 < \tan \beta < 4$  and the lightest higgs mass  $m_{h^0} < 85$  GeV. In fact, the latter provides the strongest prospect for falsifying the theory since, as shown in figure 5<sup>9</sup>,  $m_{h^0}$  up to 90 GeV should be ruled out or discovered by the completion of LEP 2. Moreover, values of  $\tan \beta$  up to 4 will be explored, which is essentially the same thing since the mass of the light Higgs is related to  $\tan \beta$  in a simple way.

A less robust signal of new physics is the relatively light top squark required. Figure 6a<sup>10</sup> shows how the sensitivity to the one-loop decay channel  $\tilde{t} \rightarrow c\tilde{\chi}_1^0$  (charm plus neutralino) depends strongly on the neutralino mass. Similarly figure 6b shows that much of the  $\mu - m_2$  plane relevant for baryogenesis is already excluded by LEP 1.5, but under the assumption of heavy sneutrinos (pair production of charginos is reduced by destructive interference from  $\tilde{\nu}$  exchange if the latter are light).

A potentially serious challenge to the naturalness of baryogenesis in the MSSM is the very large value indicated for the  $\mu$  parameter phase  $\delta$ . Values of  $\delta > 10^{-3}$  would normally be considered as incompatible with bounds on the neutron electric dipole moment, but this is only true if the lower generation squarks and the charginos are all relatively light. If the up and down squarks are much heavier than the stop and sbottom, say 1 TeV, then the EDM bound on  $\delta$  is relaxed. One might object that it looks unnatural to make a hierarchy between the first two generations and the third generation squarks. It is encouraging that such a scenario has recently been proposed for completely different reasons.<sup>11</sup>




Figure 5: LEP2 discovery potential for lightest Higgs boson in MSSM, and sensitivity to  $\tan\beta$ . Courtesy of P. Janot (ALEPH).




Figure 6: Exclusion plots for (a) neutralinos versus light stops, depending on stop mixing angle  $\theta_t$ , and (b)  $m_2$  versus  $\mu$ . Double-hatched region in (a) is excluded by D0 at Tevatron. Courtesy of J.-F. Grivaz<sup>10</sup> (ALEPH).

## 5 Summary

It appears that producing the baryon asymmetry in the MSSM is not easy; the parameter space is highly constrained. From the perspective seeking an experimentally falsifiable theory of baryogenesis, this is encouraging. The phase transition is strong enough only with a relatively light stop and higgs, and  $\tan\beta \sim 2$ . Baryon production is maximized when  $\mu \sim m_2$ , hence the Wino and Higgsino are roughly degenerate. Also the CP violation in the  $\mu$  parameter must satisfy  $\sin\delta \gtrsim 0.1$ , which indicates a large neutron EDM, in fact too large unless the lower generation squarks are much heavier than the light stop. If supersymmetry should be confirmed in the next few years and these predictions verified, it will be very tempting to believe we finally have evidence for the true origin of the baryon asymmetry of the universe.

## References

1. A.E. Nelson, D.B. Kaplan and A.G. Cohen, *Nucl. Phys. B* **373**, 453 (1992).
2. P. Huet and A.E. Nelson, *Phys. Rev. D* **53**, 4578 (1996), hep-ph/9506477; M. Carena, M. Quiros, A. Riotto, I. Vilja and C.E.M. Wagner, *Nucl. Phys. B* **503**, 404 (1997).
3. B. de Carlos and J.R. Espinosa, *Nucl. Phys. B* **503**, 24 (1997) and references therein.
4. J.M. Cline and K. Kainulainen, *Nucl. Phys. B* **482**, 73 (1996), M. Laine, *Nucl. Phys. B* **481**, 43 (1996), M. Losada, *Phys. Rev. D* **56**, 2893 (1997).
5. K. Kajantie, M. Laine, K. Rummukainen and M.E. Shaposnikov, *Nucl. Phys. B* **466**, 189 (1996), hep-lat/9510020.
6. J.M. Cline and K. Kainulainen, hep-ph/9705201 (1997), to appear in *Nucl. Phys. B*, (1998).
7. M. Joyce, T. Prokopec and N. Turok, *Phys. Rev. Lett.* **75**, 1695 (1995).
8. J.M. Cline, M. Joyce and K. Kainulainen, "Supersymmetric Electroweak Baryogenesis in the WKB Approximation," hep-ph/9708393 (1997), to appear in *Phys. Lett. B*, (1998).
9. "Physics at LEP2 (Yellow Book)," preprint CERN 96-01 (1996).
10. J.F. Grivaz, hep-ph/9709505, to appear in "Perspectives on Supersymmetry," World Scientific Publishing Co., Gordon L. Kane, Ed.
11. G. Dvali and A. Pomarol, *Phys. Rev. Lett.* **77**, 3728 (1996); A.G. Cohen, D.B. Kaplan and A.E. Nelson, *Phys. Lett. B* **388**, 588 (1996).